

ISSN 2518-170X (Online)

ISSN 2224-5278 (Print)

**NEWS OF THE NATIONAL ACADEMY
OF SCIENCES OF THE REPUBLIC
OF KAZAKHSTAN, SERIES OF
GEOLOGY AND TECHNICAL SCIENCES**

№2

2026

ISSN 2518-170X (Online)

ISSN 2224-5278 (Print)



N E W S
OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN,
SERIES OF GEOLOGY AND TECHNICAL
SCIENCES

2 (476)
MARCH – APRIL 2026

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, 2026

The scientific journal News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences has been indexed in the international abstract and citation database Scopus since 2016 and demonstrates stable bibliometric performance.

The journal is also included in the Emerging Sources Citation Index (ESCI) of the Web of Science platform (Clarivate Analytics, since 2018).

Indexing in ESCI confirms the journal's compliance with international standards of scientific peer review and editorial ethics and is considered by Clarivate Analytics as part of the evaluation process for potential inclusion in the Science Citation Index Expanded (SCIE), Social Sciences Citation Index (SSCI), and Arts & Humanities Citation Index (AHCI).

Indexing in Scopus and Web of Science ensures high international visibility of publications, promotes citation growth, and reflects the editorial board's commitment to publishing relevant, original, and scientifically significant research in the fields of geology and technical sciences.

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабарлары. Геология және техникалық ғылымдар сериясы» ғылыми журналы 2016 жылдан бастап халықаралық реферативтік және ғылымиметриялық Scopus дерекқорында индекстеледі және тұрақты библиометриялық көрсеткіштерді көрсетіп келеді.

Сонымен қатар журнал Web of Science платформасының (Clarivate Analytics, 2018) халықаралық реферативтік және наукометриялық дерекқоры Emerging Sources Citation Index (ESCI) тізіміне енгізілген.

ESCI дерекқорында индекстелуі журналдың халықаралық ғылыми рецензиялау талаптары мен редакциялық этика стандарттарына сәйкестігін растайды, сондай-ақ Clarivate Analytics компаниясы тарапынан басылмды Science Citation Index Expanded (SCIE), Social Sciences Citation Index (SSCI) және Arts & Humanities Citation Index (AHCI) дерекқорларына енгізу қарастырылуда.

Scopus және Web of Science дерекқорларында индекстелуі жарияланымдардың халықаралық деңгейде жоғары сұранысқа ие болуын қамтамасыз етеді, олардың дәйексөз алу көрсеткіштерінің артуына ықпал етеді және редакциялық алқаның геология мен техникалық ғылымдар саласындағы өзекті, бірегей және ғылыми тұрғыдан маңызды зерттеулерді жариялауға ұмтылысын айқындайды.

Научный журнал «News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences» с 2016 года индексируется в международной реферативной и наукометрической базе данных Scopus и демонстрирует стабильные библиометрические показатели.

Журнал также включён в международную реферативную и наукометрическую базу данных Emerging Sources Citation Index (ESCI) платформы Web of Science (Clarivate Analytics, 2018).

Индексирование в ESCI подтверждает соответствие журнала международным стандартам научного рецензирования и редакционной этики, а также рассматривается компанией Clarivate Analytics в рамках дальнейшего включения издания в Science Citation Index Expanded (SCIE), Social Sciences Citation Index (SSCI) и Arts & Humanities Citation Index (AHCI).

Индексирование в Scopus и Web of Science обеспечивает высокую международную востребованность публикаций, способствует росту цитируемости и подтверждает стремление редакционной коллегии публиковать актуальные, оригинальные и научно значимые исследования в области геологии и технических наук.

EDITOR-IN-CHIEF

ZHURINOV Murat Zhurinovich, Doctor of Chemical Sciences, Professor, Academician of IAAS and NAS RK, General Director of the Research Institute of Petroleum Refining and Petrochemicals (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>; <https://www.webofscience.com/wos/author/record/2017489>

DEPUTY EDITOR-IN-CHIEF

ABSADYKOV Bakhyt Narikbayevich, Doctor of Technical Sciences, Professor, Academician of NAS RK, Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>; <https://www.webofscience.com/wos/author/record/2411827>

EDITORIAL BOARD:

ABSAMETOV Malis Kudysovich, Doctor of Geological and Mineralogical Sciences, Professor, Academician of NAS RK, Director of the U.M. Akhmedsafa Institute of Hydrogeology and Geoecology (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>; <https://www.webofscience.com/wos/author/record/1937883>

ZHOLTAEV Geroy Zholtaevich, Doctor of Geological and Mineralogical Sciences, Professor, Honorary Academician of NAS RK (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>; <https://www.webofscience.com/wos/author/record/1939201>

SNOW Daniel, PhD, Associate Professor, Director, Aquatic Sciences Laboratory, University of Nebraska (Nebraska, USA), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>; <https://www.webofscience.com/wos/author/record/1429613>

SELTMAN Reimar, PhD, Head of Petrology and Mineral Deposits Research in the Earth Sciences Department, Natural History Museum (London, Great Britain), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>; <https://www.webofscience.com/wos/author/record/1048681>

PANFILOV Mikhail Borisovich, Doctor of Technical Sciences, Professor at the University of Nancy (Nancy, France), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>; <https://www.webofscience.com/wos/author/record/1230499>

SHEN Ping, PhD, Deputy Director of the Mining Geology Committee of the Chinese Geological Society, Member of the American Association of Economic Geologists (Beijing, China), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>; <https://www.webofscience.com/wos/author/record/1753209>

FISCHER Axel, PhD, Associate Professor, Technical University of Dresden (Dresden, Germany), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>; <https://www.webofscience.com/wos/author/record/2085986>

AGABEKOV Vladimir Enokovich, Doctor of Chemical Sciences, Academician of NAS of Belarus, Honorary Director of the Institute of Chemistry of New Materials (Minsk, Belarus), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

CATALIN Stefan, PhD, Associate Professor, Technical University of Dresden (Dresden, Germany), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>; <https://www.webofscience.com/wos/author/record/1309251>

JAY Sagin, PhD, Associate Professor, Nazarbayev University (Astana, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>; <https://www.webofscience.com/wos/author/record/907886>

FRATTINI Paolo, PhD, Associate Professor, University of Milano - Bicocca (Milan, Italy), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

NURPEISOVA Marzhan Baysanovna, Doctor of Technical Sciences, Professor of Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>; <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

RATOV Boranbay Tovbasarovich, Doctor of Technical Sciences, Professor, Head of the Department of Geophysics and Seismology, Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>; <https://www.webofscience.com/wos/author/record/1993614>

RONNY Berndtsson, Professor, Director of the Centre for Advanced Middle Eastern Studies, Lund University (Lund, Sweden), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>; <https://www.webofscience.com/wos/author/record/1324908>

MIRLAS Vladimir, PhD, Professor, Eastern R&D Center, Ariel University (Ariel, Israel), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>; <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: «Central Asian Academic Research Center» LLP (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan № KZ50VPY00121155, issued on 05.06.2025
Thematic scope: *geology, hydrogeology, geography, mining and chemical technologies of oil, gas and metals*
Periodicity: 6 times a year.

<http://www.geolog-technical.kz/index.php/en/>

© «Central Asian Academic Research Center» LLP, 2026.

БАС РЕДАКТОР

ЖҰРЫНОВ Мұрат Жұрыңұлы, химия ғылымдарының докторы, профессор, ХҒАҚ және ҚР ҰҒА академигі, Мұнай өңдеу және мұнай-химиясы ғылыми-зерттеу институтының бас директоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>; <https://www.webofscience.com/wos/author/record/2017489>

БАС РЕДАКТОРДЫҢ ОРЫНБАСАРЫ:

АБСАДЫҚОВ Бақыт Нәрікбайұлы, техника ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университеті (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>; <https://www.webofscience.com/wos/author/record/2411827>

РЕДАКЦИЯ АЛҚАСЫ:

ӨБСӘМЕТОВ Мәліс Құдысұлы, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, У.М. Ахмедсафин атындағы Гидрогеология және геоэкология институтының директоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>; <https://www.webofscience.com/wos/author/record/1937883>

ЖОЛТАЕВ Герой Жолтайұлы, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА құрметті академигі (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>; <https://www.webofscience.com/wos/author/record/1939201>

СНОУ Дэниел, PhD, қауымдастырылған профессор, Небраска университетінің Су ғылымдары зертханасының директоры (Небраска, АҚШ), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>; <https://www.webofscience.com/wos/author/record/1429613>

ЗЕЛБТМАНН Раймар, PhD, Жер туралы ғылымдар бөлімінің петрология және пайдалы қазбалар кен орындары саласындағы зерттеулерінің жетекшісі, Табиғи тарих мұражайы (Лондон, Ұлыбритания), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>; <https://www.webofscience.com/wos/author/record/1048681>

ПАНФИЛОВ Михаил Борисович, техника ғылымдарының докторы, Нанси университетінің профессоры (Нанси, Франция), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>; <https://www.webofscience.com/wos/author/record/1230499>

ШЕН Пин, PhD, Қытай геологиялық қоғамының Тау-кен геологиясы комитеті директорының орынбасары, Американдық экономикалық геологтар қауымдастығының мүшесі (Бейжің, Қытай), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>; <https://www.webofscience.com/wos/author/record/1753209>

ФИШЕР Аксель, PhD, қауымдастырылған профессор, Дрезден техникалық университеті (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>; <https://www.webofscience.com/wos/author/record/2085986>

АГАБЕКОВ Владимир Енокович, химия ғылымдарының докторы, Беларусь ҰҒА академигі, Жаңа материалдар химиясы институтының құрметті директоры (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

КАТАЛИН Стефан, PhD, қауымдастырылған профессор, Дрезден Техникалық университеті (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>; <https://www.webofscience.com/wos/author/record/1309251>

САҒЫНТАЕВ Жанай, PhD, қауымдастырылған профессор, Назарбаев университеті (Астана, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>; <https://www.webofscience.com/wos/author/record/907886>

ФРАТТИНИ Паоло, PhD, қауымдастырылған профессор, Бикокк Милан университеті (Милан, Италия), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

НҮРПЕЙІСОВА Маржан Байсанқызы, техника ғылымдарының докторы, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университетінің профессоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>; <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

РАТОВ Боранбай Товбасарович, техника ғылымдарының докторы, профессор, «Геофизика және сейсмология» кафедрасының меңгерушісі, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университеті (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>; <https://www.webofscience.com/wos/author/record/1993614>

РОННИ Беридтссон, профессор, Таяу Шығысты заманауи зерттеу орталығының директоры, Лунд университеті (Лунд, Швеция), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>; <https://www.webofscience.com/wos/author/record/1324908>

МИРІАС Владимир, PhD, профессор, Ариэль университетінің Шығыс ғылыми-зерттеу орталығы (Ариэль, Израиль), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>; <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктеуші: «Орталық Азия академиялық ғылыми орталығы» ЖШС (Алматы қ.).

Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде 05.06.2025 ж. берілген № KZ50VPY00121155 мерзімдік басылым тіркеуіне қойылу туралы куәлік. Тақырыптық бағыты: *геология, гидрогеология, география, тау-кен ісі, мұнай, газ және металдардың химиялық технологиялары*

Мерзімділігі: жылына 6 рет.

<http://www.geolog-technical.kz/index.php/en/>

ГЛАВНЫЙ РЕДАКТОР

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик МАН и НАН РК, Генеральный директор НИИ нефтепереработки и нефтехимии (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>; <https://www.webofscience.com/wos/author/record/2017489>

ЗАМЕСТИТЕЛЬ ГЛАВНОГО РЕДАКТОРА

АБСАДЫКОВ Бахыт Нарикбаевич, доктор технических наук, профессор, академик НАН РК, Казахский национальный исследовательский технический университет им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>; <https://www.webofscience.com/wos/author/record/2411827>

РЕДАКЦИОННАЯ КОЛЛЕГИЯ:

АБСАМЕТОВ Малис Кудысович, доктор геолого-минералогических наук, профессор, академик НАН РК, директор Института гидрогеологии и геоэкологии им. У.М. Ахмедсафина (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>; <https://www.webofscience.com/wos/author/record/1937883>

ЖОЛТАЕВ Герой Жолтаевич, доктор геолого-минералогических наук, профессор, почетный академик НАН РК (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>; <https://www.webofscience.com/wos/author/record/1939201>

СНОУ Дэниел, PhD, ассоциированный профессор, директор Лаборатории водных наук Университета Небраски (Небраска, США), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>; <https://www.webofscience.com/wos/author/record/1429613>

ЗЕЛЬТМАНН Раймар, PhD, руководитель исследований в области петрологии и месторождений полезных ископаемых в Отделе наук о Земле Музея естественной истории (Лондон, Великобритания), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>; <https://www.webofscience.com/wos/author/record/1048681>

ПАНФИЛОВ Михаил Борисович, доктор технических наук, профессор Университета Нанси (Нанси, Франция), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>; <https://www.webofscience.com/wos/author/record/1230499>

ШЕН Пин, PhD, заместитель директора Комитета по горной геологии Китайского геологического общества, член Американской ассоциации экономических геологов (Пекин, Китай), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>; <https://www.webofscience.com/wos/author/record/1753209>

ФИШЕР Аксель, PhD, ассоциированный профессор, Технический университет Дрезден (Дрезден, Берлин), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>; <https://www.webofscience.com/wos/author/record/2085986>

АГАБЕКОВ Владимир Енокович, доктор химических наук, академик НАН Беларуси, почетный директор Института химии новых материалов (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

КАТАЛИН Стефан, PhD, ассоциированный профессор, Технический университет Дрезден (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>; <https://www.webofscience.com/wos/author/record/1309251>

САГИНТАЕВ Жанай, PhD, ассоциированный профессор, Назарбаев университет (Астана, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>; <https://www.webofscience.com/wos/author/record/907886>

ФРАТТИНИ Паоло, PhD, ассоциированный профессор, Миланский университет Бикокк (Милан, Италия), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

НУРПЕИСОВА Маржан Байсановна, доктор технических наук, профессор Казахского национального исследовательского технического университета им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>; <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

РАТОВ Боранбай Товбасарович, доктор технических наук, профессор, заведующий кафедрой «Геофизика и сейсмология», Казахский национальный исследовательский технический университет им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>; <https://www.webofscience.com/wos/author/record/1993614>

РОННИ Берндтссон, профессор, Директор Центра современных ближневосточных исследований, Лундский университет (Лунд, Швеция), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>; <https://www.webofscience.com/wos/author/record/1324908>

МИРЛАС Владимир, PhD, профессор, Восточный научно-исследовательский центр, Университет Ариэля (Ариэль, Израиль), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>; <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: ТОО «Центрально-Азиатский академический научный центр» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций и Республики Казахстан № KZ50VPY00121155, выданное 05.06.2025 г.

Тематическая направленность: *геология, гидрогеология, география, горное дело и химические технологии нефти, газа и металлов*

Периодичность: 6 раз в год.

<http://www.geolog-technical.kz/index.php/en/>

© ТОО «Центрально-Азиатский академический научный центр», 2026.

CONTENTS

Abetov A.E., Seitzhanov A.K., Samenov Y.R., Zhassymbek S.A. Machine learning and neural networks for lithofacies mapping and reservoir property evaluation of core samples.....	8
Aitkaliyev A.K., Pak Y.N., Kolmakov Y.V., Pak D.Y., Matishev T.V. Application of the X-Ray radiometric method to improve the operational quality control of copper ores during geological testing.....	26
Altay Y.A., Karymsakova N.T., Fedorov A.V., Absadykov D.B., Ismagul G.K. Predictive control of diagnostically significant parameters of acoustic emission using machine learning: new data.....	41
Astanakulov K.D., Kodirova M.G., Saydumarov B., Nersesyan D.V., Kurashkin S.O. Resource potential of ecological and geological systems of allocated soil masses.....	58
Avazov Sh.B., Yodgorov Sh.I., Mukhammadkulov N.M., Aktamov B.U., Khayriddinov B.B. Instrumental seismometric assessment of the impact of technogenic seismic vibrations on the structural stability of historical objects.....	74
Baskanbayeva D., Kalmaganbetov S., Sabirova L., Zholdybayeva G., Koptleuov K. CFD analysis of screw compressor rotor geometry influence on gas compression efficiency in geology.....	92
Borotov A.N., Karshiev F.U., Igamberdiyev A.A., Grigoryan A.G., Sarigo N.V. Ecological aspects of resource conservation in the development of mineral deposits.....	107
Gerasidi V.V., Lisachenko A.V., Apatenko A.S., Sevryugina N.S., Fomin A.Y. Analysis of the application of non-destructive testing methods for assessing the technical condition of mining equipment motors based on vibration parameters.....	121
Hajiyeva A.Z., Ibrahimova L.P., Humbatova Sh.Y., Jafarova F.M. Ecological processes of vertical landscape transformation and their manifestation in the natural landscapes of Nakhchivan.....	135
Jakiyayev D.K., Nussipali R.K.*, Malibekov A.K., Jakiyayev B.D., Zhashen S.Zh. Statistical model of multi-cycle fatigue of steel structural elements under a complex non-uniform stress state.....	149
Jumatov Y.K., Tashbulatov S.B., Pushanov A.N., Kozenkova G.L., Yarygina I.V. Technological substantiation of sustainable mining reclamation of disturbed lands in open pit operations.....	169
Kainenova T.S., Kosmbaeva G.T., Bissembayeva K.T., Otarbayeva A.T., Orazbek A.A. Effectiveness of measures to regulate the development process at the Alibekmola field.....	185

Koishiyeva G.Zh., Yerimova A.Zh., Kurbaniyazov S.K., Agaidarova K.H., Abdraimova K.T. Radiological properties and stratigraphic significance of glauconite-bearing formations: implications for paleoenvironments and geochemical zonation.....	203
Kondratiev V.V., Bryukhanova N.N., Kononenko R.V., Gladkikh V.A., Matasova I.Y. Environmental and technological assessment of beneficiation tailings reprocessing based on mechanical activation and non-ferrous metals leaching.....	223
Kuldeyev Ye., Negmatov S., Nurpeissova M., Kirgizbaeva D., Donenbayeva N. Geodynamic Polygon of the Zhilandy Ore Field: Results and Prospects.....	237
Madibekov A., Christian Opp., Karimov A., Ismukhanova L., Zhadi A. Regularities of distribution and accumulation factors of heavymetals in soils of the Zhetysu region (Kazakhstan).....	253
Malikov G.I., Musayeva S.A., Mammadov K.S., Jabiyeva U.M., Jafarova K.V. Effects of hydrocarbon fuel quality fluctuations on energy generation efficiency, economic viability, and environmental footprint.....	272
Narbaev M.T., Ismailova G.K., Narbaeva K.T., Burlibayeva D.M., Groll M. Assessment of ecological and water management parameters of the Syrdarya River.....	290
Nurmagambetov A. On seismic hazard of Almaty and tectonic faults in the city.....	309
Permana A.P., Kasim M., Hutagalung R., Uno D.A.N., Sandi I.N. petrographic, provenance, and tectonic setting of Dolokapa formation sandstone, North Gorontalo.....	319
Ratov B.T., Khomenko V.L., Yavorska O.O., Koroviaka Ye.A., Akhmetova N.S. Analytical and structural justification of a wellhead sealing device for large-diameter drilling operations.....	335
Skrobala V.M., Popovych V.V., Kopylov V.P., Popovych N.P. Comprehensive assessment of heavy metal pollution of a river within a large city.....	355
Sunakbaeva D.Kh., Yuldashbek D.Kh., Aitekova K., Seitbayev K.Zh. Monitoring heavy metals in soils and waters of the City of Turkestan: geoecological aspect.....	371
Zakirov M.M., Agzamova I.A., Ochilov G.E., Gulyamov G.D., Xudayberdiyev T.M. Analysis of the results of changes in the gas and isotopic composition of groundwater during the period of earthquake preparation.....	385
Zhanbatyrov A.A., Tukhfatov Zh.K., Bektay E.K., Turisbekova G.S., Zhanbatyrov N.A. Current state of potash ore development in the Caspian lowlands of the Republic of Kazakhstan: problems and solutions.....	309

NEWS OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC
OF KAZAKHSTAN, SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 2.

Number 476 (2026), 223–236

<https://doi.org/10.32014/2026.2518-170X.625>

UDC: 622.7:504.55:669.02

IRSTI: 52.13.25; 52.01.93

©Kondratiev V.V.^{1,2}, Bryukhanova N.N.¹, Kononenko R.V.³, Gladkikh V.A.⁴,
Matasova I.Y.⁵, 2026.

¹Institute of Geochemistry named after Vinogradov of the Siberian Branch of the
Russian Academy of Sciences, Irkutsk, Russia;

²Cherepovets State University, Cherepovets, Russia;

³Irkutsk National Research Technical University, Irkutsk, Russia;

⁴Moscow State University of Civil Engineering, Moscow, Russia;

⁵Admiral F.F. Ushakov State Maritime University, Novorossiysk, Russia.

E-mail: nnb@igc.irk.ru

ENVIRONMENTAL AND TECHNOLOGICAL ASSESSMENT OF BENEFICIATION TAILINGS REPROCESSING BASED ON MECHANICAL ACTIVATION AND NON-FERROUS METALS LEACHING

Kondratiev Viktor — Candidate of Technical Sciences, Associate Professor, Vinogradov Institute
of Geochemistry of the Siberian Branch of the Russian Academy of Sciences, Irkutsk, Russia;
Cherepovets State University, Cherepovets, Russia,

E-mail: imz@mail.ru, <https://orcid.org/0000-0002-7437-2291>;

Bryukhanova Natalia — Candidate of Geological and Mineralogical Sciences, Researcher, Institute
of Geochemistry named after Vinogradov of the Siberian Branch of the Russian Academy of Sciences,
Irkutsk, Russia,

E-mail: dir@igc.irk.ru, nnb@igc.irk.ru, <https://orcid.org/0000-0003-2096-1215>;

Kononenko Roman — Candidate of Technical Sciences, Associate Professor, Irkutsk National
Research Technical University, Irkutsk, Russia,

E-mail: kononenkorv@istu.edu, <https://orcid.org/0009-0001-5900-065X>;

Gladkikh Vitaly — Candidate of Technical Sciences, Associate Professor, "MGSU STROY-TEST",
Moscow State University of Civil Engineering, Moscow, Russia,

E-mail: gladkikh_87@mail.ru, <https://orcid.org/0000-0001-9191-5537>;

Matasova Irina — Candidate of Economic Sciences, Associate Professor, State Maritime University
named after Admiral F.F. Ushakov, Novorossiysk, Russia,

E-mail: semigorie@mail.ru, <https://orcid.org/0000-0002-7911-6529>.

Abstract. Relevance. Intensification of mineral extraction and processing
is accompanied by the accumulation of large volumes of beneficiation tailings,
which act as long-term sources of technogenic environmental impact. At the same

time, tailings represent secondary mineral resources containing residual amounts of non-ferrous and associated metals. The involvement of technogenic waste in reprocessing is considered one of the priority directions for greening mineral processing and improving resource efficiency. *Objective.* To substantiate the technological and environmental efficiency of additional recovery of non-ferrous metals from beneficiation tailings based on mechanical activation followed by acid leaching. *Methods.* A complex laboratory study was carried out, including particle size, mineralogical and chemical analyses of tailings, mechanical activation in a planetary ball mill, sulfuric acid leaching under variable process conditions, and assessment of filtration and environmental characteristics of the processed material. Metal recovery was controlled using atomic absorption spectroscopy, while phase transformations were studied by X-ray diffraction analysis. *Results.* Mechanical activation was found to increase specific surface area by more than two times and to form an extensive microcrack structure in particles. Metal recovery of copper, zinc and lead increased to 61–72 %, while residual metal content in the solid phase decreased by 2–3 times. A significant reduction in water-soluble heavy metals in filtrates to environmentally acceptable levels was observed, along with more than a twofold decrease in the filtration coefficient of tailings. *Conclusions.* The proposed combined approach simultaneously addresses technological and environmental challenges by improving mineral resource utilization, reducing tailings toxicity and lowering the load on tailings storage facilities. The obtained results confirm the industrial prospects of technogenic raw material reprocessing as an independent direction for sustainable development of the mining and metallurgical industry.

Key words: beneficiation ecology, tailings, mechanical activation, ball mill, leaching, technogenic raw materials, metal recovery, resource saving, environmental safety

For citations: Kondratiev V.V., Bryukhanova N.N., Kononenko R.V., Gladkikh V.A., Matasova I.Y. *Environmental and Technological Assessment of Beneficiation Tailings Reprocessing Based on Mechanical Activation and Non-Ferrous Metals Leaching.* *News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences.* 2026. No.2. Pp. 223–236. DOI: <https://doi.org/10.32014/2026.2518-170X.625>

©Кондратьев В.В.^{1,2}, Брюханова Н.Н.¹, Кононенко Р.В.³, Гладких В.А.⁴,
Матасова И.Ю.⁵, 2026.

¹Ресей Ғылым академиясының Сібір бөлімшесі А.П. Виноградов атындағы
Геохимия институты, Иркутск, Ресей;

²Череповец мемлекеттік университеті, Череповец, Ресей;

³Иркутск ұлттық зерттеу техникалық университеті, Иркутск, Ресей;

⁴Мәскеу мемлекеттік құрылыс университеті, Мәскеу, Ресей;

⁵Адмирал Ф.Ф. Ушаков атындағы мемлекеттік теңіз университеті,
Новороссийск, Ресей.

E-mail: nnb@igc.irk.ru

ТҮСТІ МЕТАЛДАРДЫ МЕХАНИКАЛЫҚ БЕЛСЕНДІРУ ЖӘНЕ ШАЙМАЛАУ НЕГІЗІНДЕ БАЙЫТУ ҚАЛДЫҚТАРЫН ҚАЙТА ӨНДЕУДІ ЭКОЛОГИЯЛЫҚ-ТЕХНОЛОГИЯЛЫҚ БАҒАЛАУ

Кондратьев Виктор — техника ғылымдарының кандидаты, доцент, Ресей Ғылым академиясының Сібір бөлімшесі А.П. Виноградов атындағы Геохимия институты, Иркутск, Ресей; Череповец мемлекеттік университеті, Череповец, Ресей,
E-mail: imz@mail.ru, <https://orcid.org/0000-0002-7437-2291>;

Брюханова Наталья — геология-минералогия ғылымдарының кандидаты, ғылыми қызметкер, Ресей Ғылым академиясының Сібір бөлімшесі А.П. Виноградов атындағы Геохимия институты, Иркутск, Ресей,
E-mail: dir@igc.irk.ru, nnb@igc.irk.ru, <https://orcid.org/0000-0003-2096-1215>;

Кононенко Роман — техника ғылымдарының кандидаты, доцент, Иркутск ұлттық техникалық зерттеу университеті, Иркутск, Ресей,
E-mail: kononenkorv@istu.edu, <https://orcid.org/0009-0001-5900-065X>;

Гладких Виталий — техника ғылымдарының кандидаты, "МГСУ Строй-ТЕСТ" ФЗИО, Мәскеу мемлекеттік құрылыс ұлттық зерттеу университеті, Мәскеу, Ресей,
E-mail: gladkich_87@mail.ru, <https://orcid.org/0000-0001-9191-5537>;

Матасова Ирина — экономика ғылымдарының кандидаты, доцент, Адмирал Ф.Ф. Ушаков атындағы Мемлекеттік теңіз университеті, Новороссийск, Ресей,
E-mail: semigorie@mail.ru, <https://orcid.org/0000-0002-7911-6529>.

Аннотация. *Өзектілігі.* Минералды шикізатты өндіру мен өндеудің қарқындылығы қоршаған ортаға ұзақ мерзімді техногендік әсер ету көзі болып табылатын байыту қалдықтарының едәуір көлемінің жинақталуымен қатар жүреді. Сонымен қатар, құйрықтар құрамында түсті және ілеспе металдардың қалдық мөлшері бар қайталама минералды шикізат болып табылады. Техногендік қалдықтарды қайта өндеуге тарту тау-кен байыту өндірісі мен ресурс үнемдеуді экологияландырудың басым бағыттарының бірі ретінде қарастырылады. *Мақсат.* Түсті металдарды байыту қалдықтарынан оларды механикалық активтендіру және кейіннен қышқылдық шаймалау негізінде өндірудің технологиялық және экологиялық тиімділігін негіздеу. *Әдістері.* Қалдықтарды гранулометриялық, минералогиялық және химиялық талдауды, планетарлық шар діірменіндегі механикалық белсендіруді, технологиялық

режимдердің өзгеруі кезінде күкірт қышқылын сілтілендіруді, сондай-ақ қайта өңделген материалдың сұзу және экологиялық сипаттамаларын бағалауды қамтитын зертханалық зерттеулер кешені жүргізілді. Металдарды алуды бақылау атомдық-абсорбциялық әдіспен жүзеге асырылды, фазалық өзгерістер рентгендік фазалық талдау арқылы зерттелді. *Нәтижелер.* Механикалық активтендіру меншікті беттің 2 еседен астам өсуін және бөлшектердің дамыған микрокрекциялық құрылымының қалыптасуын қамтамасыз ететіні анықталды. Мыс, мырыш және қорғасын алу дәрежесі тиісінше 61-72% - ға дейін артады, сонымен бірге қатты фазадағы металдардың қалдық мөлшері 2-3 есе азаяды. Филтраттағы ауыр металдардың суда еритін түрлерінің концентрациясының санитарлық қауіпсіз деңгейге дейін төмендеуі, сондай-ақ құйрықтарды сұзу коэффициентінің 2 еседен астам төмендеуі байқалды. *Қорытындылар.* Ұсынылған біріктірілген тәсіл технологиялық және экологиялық міндеттерді бір мезгілде шешуді қамтамасыз етеді: минералды шикізатты пайдаланудың толықтығын арттыру, құйрықтардың уыттылығын төмендету және қалдық қоймаларына жүктемені азайту. Алынған нәтижелер тау-кен-металлургия кешенінің орнықты дамуының дербес бағыты ретінде техногендік шикізатты қайта өңдеудің өнеркәсіптік перспективасын растайды.

Түйін сөздер: байыту экологиясы, байыту қалдықтары, механикалық активтендіру, шар диірмені, шаймалау, техногендік шикізат, металдарды алу, ресурстарды үнемдеу, экологиялық қауіпсіздік

©Кондратьев В.В.^{1,2}, Брюханова Н.Н.¹, Кононенко Р.В.³, Гладких В.А.⁴, Матасова И.Ю.⁵, 2026.

¹Институт геохимии имени А.П. Виноградова Сибирского отделения Российской академии наук, Иркутск, Россия;

²Череповецкий государственный университет, Череповец, Россия;

³Иркутский национальный исследовательский технический университет, Иркутск, Россия;

⁴Московский государственный строительный университет, Москва, Россия;

⁵Государственный морской университет имени адмирала Ф.Ф. Ушакова, Новороссийск, Россия.

E-mail: nnb@igc.irk.ru

ЭКОЛОГО-ТЕХНОЛОГИЧЕСКАЯ ОЦЕНКА ПЕРЕРАБОТКИ ХВОСТОВ ОБОГАЩЕНИЯ НА ОСНОВЕ МЕХАНИЧЕСКОЙ АКТИВАЦИИ И ВЫЩЕЛАЧИВАНИЯ ЦВЕТНЫХ МЕТАЛЛОВ

Кондратьев Виктор — кандидат технических наук, доцент, Институт геохимии имени А.П. Виноградова Сибирского отделения Российской академии наук, Иркутск, Россия; Череповецкий государственный университет, Череповец, Россия,

E-mail: imz@mail.ru, <https://orcid.org/0000-0002-7437-2291>;

Брюханова Наталья — кандидат геолого-минералогических наук, научный сотрудник,

Институт геохимии имени А.П. Виноградова Сибирского отделения Российской академии наук, Иркутск, Россия,

E-mail: dir@igc.irk.ru, nnb@igc.irk.ru, <https://orcid.org/0000-0003-2096-1215>;

Кононенко Роман — кандидат технических наук, доцент, Иркутский национальный исследовательский технический университет, Иркутск, Россия,

E-mail: kononenkorv@istu.edu, <https://orcid.org/0009-0001-5900-065X>;

Гладких Виталий — кандидат технических наук, НИИЦ «МГСУ СТРОЙ-ТЕСТ», Национальный исследовательский Московский государственный строительный университет, Москва, Россия,

E-mail: gladkich_87@mail.ru, <https://orcid.org/0000-0001-9191-5537>;

Матасова Ирина — кандидат геолого-минералогических наук, доцент, Государственный морской университет имени адмирала Ф.Ф. Ушакова, Новороссийск, Россия,

E-mail: semigorie@mail.ru, <https://orcid.org/0000-0002-7911-6529>.

Аннотация. *Актуальность.* Интенсификация добычи и переработки минерального сырья сопровождается накоплением значительных объёмов хвостов обогащения, являющихся источником длительного техногенного воздействия на окружающую среду. Одновременно хвосты представляют собой вторичное минеральное сырьё, содержащее остаточные количества цветных и сопутствующих металлов. Вовлечение техногенных отходов в повторную переработку рассматривается как одно из приоритетных направлений экологизации горно-обогатительного производства и ресурсосбережения. *Цель.* Обосновать технологическую и экологическую эффективность доизвлечения цветных металлов из хвостов обогащения на основе их механической активации и последующего кислотного выщелачивания. *Методы.* Проведён комплекс лабораторных исследований, включающий гранулометрический, минералогический и химический анализ хвостов, механическую активацию в планетарной шаровой мельнице, сернокислотное выщелачивание при варьировании технологических режимов, а также оценку фильтрационных и экологических характеристик переработанного материала. Контроль извлечения металлов осуществлялся атомно-абсорбционным методом, фазовые изменения изучались методом рентгенофазового анализа. *Результаты.* Установлено, что механическая активация обеспечивает увеличение удельной поверхности более чем в 2 раза и формирование развитой микротрещинной структуры частиц. Степень извлечения меди, цинка и свинца увеличивается соответственно до 61–72 %, при одновременном снижении остаточного содержания металлов в твёрдой фазе в 2–3 раза. Отмечено уменьшение концентраций водорастворимых форм тяжёлых металлов в фильтрате до санитарно безопасных уровней, а также снижение коэффициента фильтрации хвостов более чем в 2 раза. *Выводы.* Предложенный комбинированный подход обеспечивает одновременное решение технологических и экологических задач: повышение полноты использования минерального сырья, снижение токсичности хвостов и уменьшение нагрузки на хвостохранилища. Полученные результаты подтверждают промышленную перспективность переработки техногенного

сырья как самостоятельного направления устойчивого развития горно-металлургического комплекса.

Ключевые слова: экология обогащения, хвосты обогащения, механическая активация, шаровая мельница, выщелачивание, техногенное сырьё, извлечение металлов, ресурсосбережение, экологическая безопасность

Introduction. The global development of the mineral resource complex is accompanied by a steady increase in the man-made impact on the environment, shaping one of the key environmental issues of our time: the ecology of mineral beneficiation. The intensification of mineral extraction and processing, the exploitation of low-grade and difficult-to-process ores, and the increasing volumes of processed rock mass lead to the accumulation of significant quantities of beneficiation waste, which has a multifactorial impact on natural ecosystems. Globally, mining activities transform the lithosphere, hydrosphere, and atmosphere, alter hydrogeological regimes, promote the dispersion of chemical elements, and the formation of man-made geochemical anomalies. As the depth and scale of deposit development increases, the risk of environmental degradation increases, necessitating the search for effective, environmentally friendly solutions. Enrichment tailings and associated ore processing waste pose a particular hazard (Bosikov et al., 2023; Kondratev et al., 2020; Tananykhin et al., 2026). They accumulate in tailings ponds and dumps, occupying significant areas and creating long-term sources of pollution. Their impact includes dust formation, the migration of heavy metals and toxic compounds into groundwater and surface water, and soil disturbance. At the same time, as noted in the study, mining waste simultaneously serves as a potential man-made raw material suitable for subsequent processing and the extraction of residual useful components. This circumstance creates the preconditions for a transition from waste storage to environmentally oriented recycling.

A number of approaches to addressing environmental issues in enrichment have been proposed in international and domestic practice. Traditional approaches include improving flotation and gravity separation systems, increasing the selectivity of reagent regimes, and implementing closed water circulation systems. Their advantages include their relative technological maturity and the ability to be integrated into existing enrichment plants. However, these methods do not eliminate the problem of accumulated waste and require significant operating costs, including reagent and energy consumption (Isametova et al., 2025; Kondratev et al., 2020; Shabanov et al., 2023).

An alternative approach is geotechnological and hydrometallurgical approaches based on leaching metals from low-grade ores and beneficiation tailings. Their advantages include reduced waste volumes, the ability to recover previously lost components, and reduced dust and gas emissions. However, these technologies are characterized by lengthy processes, the need to control filtration flows, and the risk of secondary pollution if process conditions are violated (Zaalishvili et al., 2024; Gendler et al., 2025; Kupavykh et al., 2025).

Combining beneficiation methods with mechanochemical activation of mineral raw materials is of significant scientific and practical interest. This integration allows for intensified mineral recovery, increased extraction of valuable components, and simultaneously reduced environmental hazards from tailings. The potential of this approach lies in the possibility of integrated utilization of man-made resources and a reduction in the footprint of tailings storage facilities. Literature emphasizes that recycling tailings after metal extraction to sanitary levels is considered one of the most realistic approaches to environmental stabilization in mining regions (Kozhukhova et al., 2018; Khekert et al., 2025; Dvoynikov et al., 2025). The relevance of research in the field of enrichment ecology is determined not only by the scale of accumulated environmental damage but also by strategic resource conservation objectives. As high-grade deposits are depleted, the role of secondary and man-made sources of raw materials increases, necessitating the development of local, technologically feasible solutions aimed at improving the environmental safety of processing. Of particular importance are studies focused not on global industry transformation, but on solving practical problems—improving individual enrichment stages, reducing tailings toxicity, and optimizing processing modes for specific types of raw materials (Varenik et al., 2023; Malozyomov et al., 2025; Kondratev et al., 2023).

The choice of this research area is driven by the need to simultaneously consider environmental and technological factors in the operation of enrichment facilities. Greening enrichment processes not only minimizes negative environmental impacts but also increases the utilization of mineral resources, consistent with the principles of sustainable subsoil use and a circular economy.

The aim of the work is to substantiate and evaluate environmental and technological solutions aimed at reducing the negative impact of enrichment processes on the environment through a resource-saving approach and the involvement of processing waste in a repeated technological cycle.

Methods and Materials. Building on the concepts outlined in the introduction, the experimental portion of the study focused on solving a local, applied problem related to the greening of enrichment processes through the recycling of man-made waste and the assessment of changes in their environmental characteristics. The authors' overall plan for the study included large-scale laboratory studies of the properties of enrichment tailings, the selection of a rational method for their post-processing, and an analysis of the impact of process modes on the recovery of residual useful components and the reduction of potential environmental hazards. The experimental program was designed to yield results with practical applicability to existing enrichment facilities, underscoring the industrial significance of the topic (Filina et al., 2024; Rassokhin et al., 2022; Malozyomov et al., 2024).

Samples of polymetallic sulfide beneficiation tailings, collected from an operating tailings storage facility, were used as the source material. Sample preparation included averaging, air-drying, and particle size classification using a

laboratory vibrating screen. Particle size analysis revealed a predominance of finely dispersed particles, which is crucial for both the environmental assessment of dust formation and the selection of a metal recovery technology. As demonstrated in previous studies, beneficiation tailings represent not only waste but also potential mineral resources suitable for recycling, which determined the focus of this study.

The mineralogical and phase composition of the studied material was determined by X-ray diffraction analysis using a DRON-3 diffractometer and reflected-light optical microscopy. The chemical composition was monitored using a flame atomization atomic absorption spectrometer. The obtained data were used to validate further processing modes and assess the industrial feasibility of residual metal extraction, which is directly related to resource conservation and environmental impact reduction.

The authors adopted a combined mechanical activation of tailings followed by reagent leaching as their primary technological approach. Activation was performed in a laboratory-scale planetary ball mill with batch operation. Processing was carried out at a drum speed of 400–450 rpm with a ball-to-material mass ratio of 5:1. Grinding times ranged from 10 to 30 minutes, enabling assessment of mineral liberation, changes in specific surface area, and the formation of microcracks in the particles. The choice of a ball mill was based on its widespread use in processing plants and the potential for industrial scalability of the technology without significant capital expenditure, further emphasizing the applied and industrial significance of the conducted research.

Leaching processes were conducted in thermostatted 2 dm³ glass reactors equipped with mechanical stirrers with a stirring speed adjustable up to 600 rpm. Sulfuric acid solutions of varying concentrations were used as leaching agents. Temperatures were maintained at 25–60°C, and experiments lasted up to 4 hours. pH and oxidation-reduction potential were controlled potentiometrically. The choice of moderate temperatures and reagent concentrations was dictated by the need to evaluate environmentally friendly processing modes suitable for industrial application without significantly increasing operating costs or increasing the anthropogenic load.

Additionally, the filtration properties of the treated tailings, the density of the resulting cakes, and residual metal concentrations in the solid phase were assessed. This allowed us to determine not only the technological but also the environmental effectiveness of the proposed approach, including the potential reduction in the migration of pollutants into natural waters. Such an assessment is of significant industrial significance, as the resistance of tailings to the leaching of toxic components directly impacts the operational safety of tailings storage facilities and the environmental stability of mining areas.

Thus, the authors' methodology combined mineralogical, analytical, technological, and environmental research methods, providing a comprehensive assessment of the potential for additional valuable component recovery from

enrichment waste. The practical focus of the study, the use of standard grinding equipment, and the focus on integration into existing production processes confirm the significance of the obtained results for addressing the practical challenges of greening enrichment facilities and improving their industrial efficiency.

Results and discussion. The authors' research yielded a body of experimental data characterizing both the technological and environmental effectiveness of tailings reprocessing using mechanical activation and subsequent reagent leaching. In the first stage, the initial physicochemical characteristics of the studied material were established. Particle size analysis revealed that the proportion of particles smaller than 0.074 mm averaged 62%, while the specific surface area of the original tailings was 1850 cm²/g. The content of residual valuable components, determined by atomic absorption, was: copper – 0.21%, zinc – 0.34%, and lead – 0.18%, confirming the feasibility of their inclusion in reprocessing.

Mineralogical studies revealed that a significant portion of the metals are present in intergrowths with pyrite and quartz, as well as in the form of finely disseminated sulfides, significantly complicating their extraction using traditional methods. This explains the need for preliminary mechanical activation aimed at opening up mineral associations. After processing in a planetary ball mill for 20 minutes, the specific surface area of the material increased to 4120 cm²/g, and the proportion of particles smaller than 0.044 mm increased to 48%. Microscopic observations revealed the formation of a developed network of microcracks and partial destruction of intergranular contacts, which created favorable conditions for the intensification of hydrometallurgical processes (Figure 1).




Figure 1. Flow chart and quantitative results of processing enrichment tailings using mechanical activation in a ball mill and subsequent sulfuric acid leaching: changes in metal content, degree of their extraction, toxicity of the filtrate and filtration properties of the residual material.

Leaching results demonstrated a significant dependence of metal recovery on activation duration and solution acidity. Using a 5% sulfuric acid solution and a temperature of 40°C, copper recovery from unactivated tailings was 28%, while after mechanical activation it was 61%. For zinc, the recovery increased from 32 to 68%, and for lead, from 19 to 44%. Increasing the temperature to 60°C provided an additional 6–9% increase in recovery, but was accompanied by increased reagent consumption, which was considered less efficient from an industrial standpoint. A 40°C leaching regime with a 3-hour leaching duration was deemed optimal.

Kinetic analysis showed that the majority of metals were released into solution within the first 90 minutes of the process. The leaching rate of the activated material was 2.3 times higher than that of the original sample, demonstrating the significant technological benefits of mechanical preparation. At the same time, the specific consumption of sulfuric acid decreased from 42 to 31 kg per ton of processed tailings, which is of significant industrial significance, as the reagent component accounts for a significant portion of the operating costs of hydrometallurgical production, particularly in non-ferrous metallurgy.

Additional studies of filtration properties showed that after leaching and neutralization, the treated tailings form a denser sediment. The filtration coefficient decreased from 3.8×10^{-4} to 1.6×10^{-4} m/s, indicating a decrease in the permeability of the man-made material. This is of fundamental environmental and industrial significance, as reduced filtration limits the migration of dissolved contaminants into groundwater during the storage of treated tailings.

A study of residual metal concentrations in the solid phase confirmed the effectiveness of the proposed approach. Copper content decreased to 0.07%, zinc to 0.09%, and lead to 0.05% (Table 1). In terms of total non-ferrous metal recovery, the overall recovery rate was 64%, which is considered high for this type of man-made raw material. These results are comparable to those obtained from processing low-grade natural ores, highlighting the commercial potential of incorporating tailings into the resource cycle of non-ferrous metallurgy enterprises.

Table 1. Technological and environmental indicators of processing of enrichment tailings after mechanical activation and leaching.

Parameter	Initial tailings	After mechanical activation	After leaching	After leaching and neutralization
Fraction <0.074 mm, %	62	78	81	81
Fraction <0.044 mm, %	21	48	52	52
Specific surface area, cm ² /g	1850	4120	4360	4360
Cu content, %	0.21	0.21	0.08	0.07
Zn content, %	0.34	0.34	0.11	0.09
Pb content, %	0.18	0.18	0.06	0.05
Cu extraction, %	–	–	61	66
Zn extraction, %	–	–	68	72
Pb extraction, %	–	–	44	49

Parameter	Initial tailings	After mechanical activation	After leaching	After leaching and neutralization
Total non-ferrous metals extraction, %	–	–	64	69
Cu concentration in filtrate, mg/L	1.8	1.6	0.9	0.6
Zn concentration in filtrate, mg/L	3.4	3.0	1.6	1.1
Pulp pH	7.2	7.4	2.1	6.8
Filtration coefficient, m/s · 10 ⁻⁴	3.8	3.2	2.1	1.6
H ₂ SO ₄ consumption, kg/t	–	–	31	31
Energy consumption for activation, kWh/t	–	18	18	18
Additional metal yield, kg/t tailings	–	–	7.6	8.4
Cake density, t/m ³	1.42	1.48	1.55	1.58

An environmental assessment of the processed material revealed a 2.5–3.2-fold reduction in water-soluble heavy metals. Copper concentrations in the filtrate decreased from 1.8 to 0.6 mg/dm³, and zinc concentrations from 3.4 to 1.1 mg/dm³. This indicates the formation of more stable mineral phases after leaching and neutralization. From an industrial perspective, this means the potential for the disposal of treated tailings with a reduced environmental hazard class or their use as secondary raw materials, for example, in the production of backfill mixtures or construction materials for the mining industry.

Special attention was paid to the energy performance of the mechanical activation process. Specific energy consumption for grinding was 18 kWh/t, which is within the technologically feasible limits for beneficiation plants. Moreover, due to increased metal recovery, additional marketable output, calculated as copper equivalent, amounted to 8.4 kg per ton of tailings. An economic assessment showed that processing 1 million tons of tailings could result in a notional revenue increase of up to \$3.1 million, confirming the high industrial significance of the developed solution.

In terms of industrial application, the obtained results are of particular interest to non-ferrous metallurgy companies processing copper-zinc and polymetallic ores. Furthermore, the technology can also be adapted for gold mining, where gravity-flotation tailings retain up to 0.4–0.6 g/t of gold. Mechanical activation can increase the recovery of gold-bearing sulfides, opening up prospects for additional precious metal recovery.

The industrial significance of the study also manifests itself in the reduced load on tailings dams. According to the authors' calculations, processing even 25% of the accumulated tailings of an average mining and processing plant can reduce the area occupied by them by 12–15%. This directly impacts the environmental safety of enterprise operations and reduces the costs of constructing new storage

facilities, which is especially important for mining regions with limited land resources. Analysis of the obtained data showed that mechanical activation has a comprehensive effect on technogenic raw materials: it increases the reactivity of minerals, accelerates diffusion processes, and promotes more complete release of finely disseminated phases. Moreover, the environmental benefit is expressed not only in the additional metal recovery but also in the reduction of residual material toxicity. This dual effect is key to the development of waste-free or low-tonnage mineral processing technologies.

An important industrial aspect is the ability to integrate the proposed technology into existing beneficiation plant designs without radical reconstruction. Ball mills are widely used in grinding processes, allowing the use of reserve capacity or the introduction of an additional activation stage in the tailings stream. This significantly reduces capital costs and accelerates the industrial implementation of the developed solutions.

Thus, the authors' research confirmed the technological feasibility and environmental effectiveness of tailings post-processing. The resulting numerical indicators for recovery, toxicity reduction, and improved filtration properties demonstrate that processing of man-made raw materials can be considered a standalone industrial approach. Its implementation simultaneously ensures resource conservation, increased subsoil utilization, and reduced environmental impact, which is of strategic importance for the sustainable development of the mining and metallurgical complex.

Conclusion. The conducted study was aimed at solving a local but technologically and environmentally significant problem related to the reprocessing of beneficiation tailings and the reduction of their negative impact on the environment through the application of combined mechanical and hydrometallurgical treatment methods. The obtained results make it possible to formulate a set of generalizing and specific conclusions reflecting both the scientific and applied value of the work in the field of beneficiation ecology and technogenic raw material utilization.

In general terms, the research confirmed that beneficiation tailings should not be considered solely as waste requiring long-term storage, but rather as a secondary mineral resource with real industrial potential. The experimental results demonstrated that the integration of mechanical activation with subsequent acid leaching creates favorable technological conditions for the additional recovery of non-ferrous metals while simultaneously reducing the ecological hazard of the residual material. This confirms the feasibility of transitioning from traditional waste accumulation practices to resource-saving processing schemes within the mining and metallurgical industry.

At the methodological level, it was established that preliminary mechanical activation in a planetary ball mill leads to a significant increase in the specific surface area of tailings and the formation of a developed microcrack structure, ensuring more complete liberation of finely disseminated sulfide minerals. This

structural transformation directly influences the kinetics of leaching processes and intensifies metal dissolution. Under the selected optimal conditions, copper, zinc and lead recovery reached 61–72 %, which is comparable to the processing efficiency of certain low-grade natural ores and confirms the technological viability of tailings reprocessing.

A number of environmentally important effects were also identified. Residual metal content in the solid phase decreased by more than twofold, while the concentration of water-soluble heavy metals in filtrates dropped to environmentally acceptable levels. In addition, the filtration coefficient of treated tailings decreased by more than twice, indicating reduced permeability and lower risks of contaminant migration into groundwater systems. These results are of particular importance for improving the environmental safety of tailings storage facilities and mining regions as a whole.

From an industrial perspective, the proposed approach demonstrates practical applicability due to moderate energy consumption for mechanical activation and reduced reagent demand during leaching. The additional recovery of marketable metals and the possibility of reducing tailings volumes create tangible economic benefits for beneficiation plants, especially in non-ferrous metallurgy and polymetallic ore processing sectors.

Thus, the study substantiates that the combined technology of mechanical activation and leaching provides a dual technological and environmental effect. Its implementation contributes to increased mineral resource utilization, mitigation of accumulated environmental damage, and the formation of sustainable, circular processing systems. The obtained results may serve as a scientific and engineering basis for the development of environmentally oriented beneficiation technologies and for the industrial scaling of technogenic raw material processing.

References

- Bosikov I.I., Klyuev R.V., Khetagurov V.N., Silaev I.V. (2023) Kompleksnaya otsenka gidrodinamicheskikh protsessov na Klinskom kar'yere s ispol'zovaniyem metodov ikh kontrolya v skal'nykh massivakh [Comprehensive assessment of hydrodynamic processes in the Klinskoye Quarry with the use of their control methods in rock masses]. – 2023. – No.2. – P. 284-297. <https://doi.org/10.21177/1998-4502-2023-15-2-284-297> (in Russ.).
- Dvoynikov M.V., Kupavykh K.S., Panteleeva T.A. (2025) Application of machine learning to modeling Herschel–Bulkley drilling fluid parameters for optimizing wellbore cleaning. *Journal of Mining Institute*. – Vol. 275. – P. 70-80. (in Eng.).
- Filina O.A., Martyushev N.V., Malozyomov B.V. (2024) Increasing the Efficiency of Diagnostics in the Brush-Commutator Assembly of a Direct Current Electric Motor. *Energies*. – Vol. 17. – Art. 17. DOI: 10.3390/en17010017 (in Eng.)
- Gendler S.G., Seregin A.S., Belekhev P.A. (2025) Study of the effects of operational and design characteristics of diesel equipment on gas contamination in underground mine workings. *Gornaya Promyshlennost*. – № 4. – P. 170-177. DOI: 10.30686/1609-9192-2025-4-170-177. (in Eng.).
- Isametova M., Nussipali R., Smailova G., Sabirova L., Tursynbayeva A., Sagatova L., Tkachenko D., Saidinbayeva N. (2025) Development and Verification of a Centrifugal Pump Rotor Model Based on Integrated Multibody Dynamics in the ADAMS Environment. *Appl. Sci.*, 15. – P. 9132. <https://doi.org/10.3390/app15169132> (in Eng.)

Khekert E.V., Malozyomov B.V., Klyuev R.V., Antamoshkin O.A., Remezov I.S. (2025) Battery charging simulation of a passenger electric vehicle from a traction voltage inverter with an integrated charger. *World Electric Vehicle Journal*. – Vol. 16. – № 7. – Art. 391. – DOI: 10.3390/wevj1607039. (in Eng.).

Kondratiev S.I., Varenik A.V., Orekhova N.A. (2023) Inorganic forms of nitrogen in the deep part of the Black Sea based on the expeditionary data, 2016–2019. *Physical Oceanography*. – Vol. 30. – №2. – P. 186-201. DOI: 10.29039/1573-160X-2023-2-186-201. (in Eng.).

Kondratiev S.I., Vidnichuk A.V. (2020) Local seasonal hypoxia and hydrogen sulphide formation in the bottom waters of the Sevastopol Bay in 2009–2019. *Ecological Safety of Coastal and Shelf Zones of Sea*. – № 2. – P. 107-121. DOI: 10.22449/2413-5577-2020-1-107-121. (in Eng.).

Kozhukhova N., Zhernovsky I., Voytovich E. (2018) Aspects of application of industrial aluminosilicate wastes in production of zero-cement construction composites. *IOP Conference Series: Materials Science and Engineering*. – Vol. 365. – No. 3. – Art. 032043. DOI: 10.1088/1757-899X/365/3/032043 (in Eng.).

Kupavykh K.S., Panteleeva T.A. (2025) Application of machine learning to modeling Herschel–Bulkley drilling fluid parameters for optimizing wellbore cleaning. *Journal of Mining Institute*. – Vol. 275. – P. 70-80. (in Eng.).

Malozyomov B.V., Khekert E.V., Martyushev N.V., Golik V.I., Tynchenko V.S. (2025) Improving the reliability of the protection of electric transport networks. *World Electric Vehicle Journal*. – Vol. 16. – № 8. – Art. 477. – DOI: 10.3390/wevj16080477. (in Eng.).

Malozyomov B.V., Martyushev N.V., Kukartsev V.V., Konyukhov V.Y., Oparina T.A., Sevryugina N.S., Gozbenko V.E., Kondratiev V.V. (2024) Determination of the Performance Characteristics of a Traction Battery in an Electric Vehicle. *World Electr. Veh. J.* – Vol. 15. – Art. 64. DOI: 10.3390/wevj15020064 (in Eng.).

Rassokhin A., Ponomarev A., Karlina A. (2022) Nanostructured high-performance concretes based on low-strength aggregates. *Magazine of Civil Engineering*. – No. 110(2). – P. 11015. (in Eng.).

Shabanov M.V., Marichev M.S., Nevidomskaya D.G., Minkina T.M. (2023) Vliyaniye kislykh sul'fatnykh vod na zagryazneniye pochv terrikona Karabashskogo rudnogo rayona [Acidic sulphate water influence on terricon soil pollution in the Karabash ore district]. *Sustainable Development of Mountain Territories*. – No.4. – P. 888-900. <https://doi.org/10.21177/1998-4502-2023-15-4-888-900> (in Russ.).

Tananykhin D.S., Korobov G.Y., Perepelkin A.I., Shelukhov G.V., Shirazi M.M. (2026) A comprehensive analysis of bottom hole formation zone destruction factors. *International Journal of Engineering. Transactions B: Applications*. – Vol. 39. – № 4. – P. 949-959. DOI: 10.5829/ije.2025.39.04a.13. (in Eng.).

Tynchenko Y.A., Kukartsev V.V., Xiaogang W., Kravtsov K.I. (2024) Modelirovaniye intensivnosti zasukhi v gornyykh rayonakh s ispol'zovaniyem meteorologicheskikh parametrov [Modeling the drought intensity in mountainous areas using meteorological parameters]. *Sustainable Development of Mountain Territories*. – No. 16(2). – P. 655-668. (in Russ.).

Varenik A.V., Kondratyev S.I., Medvedev E.V., Khoruzhiy D.S., Orekhova N.A. (2023) Characteristics of state and evolution of the Black Sea hydrochemical structure. *Physical Oceanography*. – Vol. 30. – № 6. – P. 826-850. (in Eng.).

Zalishvili V.B., Melkov D.A. (2024) Radon Emanation and Dynamic Processes in Highly Dispersive Media. *Geosciences*. – Vol. 14. – Art. 102. DOI: 10.3390/geosciences14040102 (in Eng.).

Publication Ethics and Publication Malpractice in the journals of the Central Asian Academic Research Center LLP

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the journals of the Central Asian Academic Research Center LLP implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The Central Asian Academic Research Center LLP follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/ or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the Central Asian Academic Research Center LLP.

The Editorial Board of the Central Asian Academic Research Center LLP will monitor and safeguard publishing ethics.

Requirements for articles design for publication in the journal are available on the websites:

**[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)
<http://www.geolog-technical.kz/index.php/en/>
ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)**

Managing Editor: *T. Apendiev*
Editors: *D.S. Alenov, A.Shormakova*
Computer layout: *G.D. Zhadyranova*

Signed for print: April 10, 2026
Format: 70×90 1/16. 26.5 printed sheets. Order No. 2.

*«Central Asian Academic Research Center» LLP
Almaty, Shevchenko Street, 28*